



# The Great Plains Laboratory, Inc.

William Shaw, Ph.D Director

11813 W. 77th Street, Lenexa, KS 66214

(913) 341-8949

Fax (913) 341-6207

**GPL-MYCOTOX**

Requisition #	717017	Physician Name	NO PHYSICIAN
Patient Name	Thomas Strohacker	Date of Collection	8/7/2019
Patient BirthDate	7/25/1965	Time of Collection	9:00 AM
Sex	M	Print Date	8/15/2019

## Mycotox Profile



Testing performed by The Great Plains Laboratory, Inc., Lenexa, Kansas. The Great Plains Laboratory has developed and determined the performance characteristics of this test. The test has not been evaluated by the U.S. Food and Drug Administration. The FDA does not currently regulate such testing.





Patient Name: Thomas Strohhacker Requisition #: 717017

**GPL-MYCOTOX**

**Ochratoxin:** Ochratoxin A (OTA) is a nephrotoxic, immunotoxic, and carcinogenic mycotoxin. This chemical is produced by molds in the *Aspergillus* and *Penicillium* families. Exposure is done primarily through water damaged buildings. Minimal exposure can occur through contaminated foods such as cereals, grape juices, dairy, spices, wine, dried vine fruit, and coffee. Exposure to OTA can also come from inhalation exposure in water-damaged buildings. OTA can lead to kidney disease and adverse neurological effects. Studies have shown that OTA can lead to significant oxidative damage to multiple brain regions and is highly nephrotoxic. Dopamine levels in the brain of mice have been shown to be decreased after exposure to OTA. Some studies have hypothesized that OTA may contribute to the development of neurodegenerative diseases such as Alzheimer's and Parkinson's. Treatment should be aimed at removing the source of exposure. Agents such as oral cholestyramine, charcoal, and phenylalanine can help prevent the absorption of these toxins from food. Antioxidants such as vitamins A, E, C, NAC, rosmarinic acid, and liposomal glutathione alone or in combination have been shown to mitigate the oxidative effects of the toxin. Bentonite or zeolite clay is reported to reduce the absorption of multiple mycotoxins found in food, including OTA. Studies have also shown that OTA is present in sweat, which supports the use of sauna as a treatment to increase the excretion of OTA. (PMID 17195275, 16621780, 16293235, 27521635, 22069626, 24792326, 22253638, 16140385, 2467220, 16844142, 19148691, 22069658, 16019795, 18286403, 15781206, 11439224, 17092826, 32710148)

**Dihydrocitrinone:** Dihydrocitrinone is a metabolite of Citrinin(CTN), which is a mycotoxin that is produced by the mold species *Aspergillus*, *Penicillium*, and *Monascus*. CTN exposure can lead to nephropathy, because of its ability to increase permeability of mitochondrial membranes in the kidneys. The three most common exposure routes are through ingestion, inhalation, and skin contact. CTN has been shown to be carcinogenic in rat studies. Multiple studies have linked CTN exposure to a suppression of the immune response. PMID: 11567776, 24048364, 10788357