

September 30,2019

Dr. Briemberg

On October 6, 2019 my voice will be silenced forever. A result of a corrupt medical system which gives doses of toxins to regulate cholesterol. Without an understanding of the drug, cardiologists are only looking at a cholesterol test, without concern about side effects.

Statin drugs have 2 billion in sales in Canada 18 billion in the USA annually and growing; the drug is a monster worldwide, where greed overshadows common sense. With unlimited resources for marketing, the drug is marketed as safe, with few side effects. The drug companies don't mention the use of mycotoxins in the drug, so people will not connect their failing health to the drug.

I have gone to great efforts to test the drug and my body. In an industry where labs are owned by medical corporations, I ran into many road blocks. Two tests on 80 mg dose came back with Ochratoxin A and Citrinin with 39ug/g (that is 1 billionth of a gram per gram) the maximum tolerance to humans of these toxins as set by international grain boards are 15 ug/g (not intended for long exposure). Doses of Lipitor contain the following ,10 mg =5 ug/g, 20 mg = 10 ug/g , 40 mg = 19 ug/g and 80 mg = 39 ug/g. There is a reason why 80 mg is not recommended, but even 40 mg is being brought into question on being potentially harmful. My urine test in June 2019 showed high levels of toxins remaining in my body (over a year and a half after stopping the drug). This explains the continued breakdown.

In June, I sent you several papers including stats from 2017 of 1500 reported cases of statin users getting ALS. I believe that number is far higher, but no one is paying attention. Dr. Briemberg, you have the power to make a difference to so many. Look into your files; find out how many ALS patients have been on a statin.

The increase of ALS cases over the last 30 years seems to mirror the sale of the drug since 1987.

I will be posting my data on line before I go on Sunday. I am disappointed that I couldn't figure this out sooner. Mycotoxin poisoning can be stopped if people are aware of the hazards.

I have met other ALS patients who have been on statins. One caregiver lost her mother in 2009 , the doctors didn't diagnosis ALS until her last weeks of life. She was on Lipitor.

Start a dialogue, and look into your files to see how many people were on statin before their ALS diagnosis. God knows I have gone as far as I could go, I pray you are more ethical than the other doctors I have met. A simple mycotoxin test early enough could allow for detoxing before damage is too severe. Because mold exposure could happen through food or environmental exposure anywhere, you could send one urine test of a patient who hasn't been on a statin and rule out my theory that mycotoxin poisoning being common through all ALS. The thousands spent in testing, care and research for ALS. A few hundred dollars more on a mycotoxin test couldn't hurt and may open a new line of reasoning.

Check out the website my friends are building at tomstrohhacker.com

Make a difference

Thomas Martin Strohhacker